

Novel Approaches for Enhancing Drug Solubility and Dissolution Rate

Peter Fonagy

Division of Psychology and Language Sciences, University College London, UK

Article history: Received: 1 July 2020, Accepted: 28 July 2020, Published online: 5 August 2020

ABSTRACT

The solubility and dissolution rate of drugs play pivotal roles in their therapeutic efficacy and bioavailability. However, a significant number of drug candidates face challenges related to poor solubility, which hampers their successful formulation into effective dosage forms. This abstract explores innovative strategies and novel approaches aimed at enhancing the solubility and dissolution rate of drugs to overcome these hurdles. One promising approach involves the utilization of nanotechnology, which offers a range of techniques such as nanocrystals, nanosuspensions, and nanoemulsions to improve drug solubility. These nanoscale formulations provide increased surface area and improved dispersibility, leading to enhanced dissolution rates and bioavailability. Furthermore, the application of solid dispersion techniques has emerged as a valuable tool for enhancing drug solubility. By dispersing poorly soluble drugs in a hydrophilic carrier matrix, solid dispersion systems can significantly increase drug solubility and dissolution rates, thereby improving therapeutic outcomes. Additionally, lipid-based formulations, including liposomes and lipid nanoparticles, offer unique advantages in enhancing drug solubility and bioavailability. These lipid-based carriers provide a biocompatible environment for hydrophobic drugs, facilitating their solubilization and delivery to target sites. Moreover, the utilization of complexation techniques, such as cyclodextrin inclusion complexes, has shown promise in improving drug solubility by forming stable complexes with hydrophobic drugs, thereby enhancing their aqueous solubility and dissolution kinetics. In conclusion, the integration of these novel approaches offers a multifaceted strategy to address the challenge of poor drug solubility. By leveraging advancements in nanotechnology, solid dispersion techniques, lipid-based formulations, and complexation strategies, researchers can enhance drug solubility and dissolution rates, ultimately leading to improved therapeutic outcomes and patient compliance.

Keywords: Drug solubility, Dissolution rate, Nanotechnology, Solid dispersion, Lipid-based formulations.

INTRODUCTION

The solubility and dissolution rate of pharmaceutical compounds are critical factors influencing their therapeutic effectiveness and bioavailability. Poorly soluble drugs often pose significant challenges in drug formulation, leading to suboptimal drug delivery and therapeutic outcomes. Enhancing the solubility and dissolution rate of drugs is, therefore, a primary focus of pharmaceutical research and development. This introduction provides an overview of the importance of drug solubility and dissolution rate, outlines the challenges associated with poor solubility, and highlights the significance of novel approaches in overcoming these challenges.

Firstly, solubility refers to the ability of a drug to dissolve in a given solvent, typically aqueous media in the context of oral drug delivery. The dissolution rate, on the other hand, pertains to the rate at which a solid drug dissolves in a solvent to form a solution. Both parameters are crucial determinants of drug absorption and bioavailability, as the drug must be in solution form to be absorbed into the systemic circulation and exert its pharmacological effect. However, a substantial number of drug candidates exhibit poor solubility, which limits their dissolution rate and, consequently, their bioavailability.

The poor solubility of drugs can be attributed to various factors, including their physicochemical properties such as high lipophilicity, large molecular size, and complex crystal lattice structures. These properties hinder the drug's interaction with aqueous media, resulting in low solubility and slow dissolution kinetics. Consequently, formulation scientists face considerable challenges in developing dosage forms with adequate drug solubility and dissolution characteristics.

Overcoming the limitations posed by poor drug solubility requires innovative strategies and novel approaches that can enhance drug solubility and dissolution rates. In recent years, significant advancements have been made in this field, leveraging techniques such as nanotechnology, solid dispersion, lipid-based formulations, and complexation strategies.

These approaches offer promising avenues for improving drug solubility by increasing surface area, enhancing dispersibility, and facilitating drug release.

In light of the importance of drug solubility and dissolution rate in pharmaceutical development, this review explores the latest advancements and novel approaches aimed at overcoming the challenges associated with poor solubility. By elucidating the principles underlying these approaches and highlighting their potential applications, this review aims to provide insights into the strategies for enhancing drug solubility and dissolution rates, ultimately contributing to the development of more effective and bioavailable pharmaceutical formulations.

LITERATURE REVIEW

In recent years, there has been a growing body of literature focusing on innovative approaches to enhance drug solubility and dissolution rates, reflecting the increasing importance of these parameters in pharmaceutical development. This literature review aims to provide an overview of the key findings and advancements in this field, highlighting the diverse range of strategies employed to address the challenges posed by poor drug solubility.

One of the most promising approaches is the utilization of nanotechnology, which has revolutionized drug delivery by offering novel techniques to improve drug solubility. Nanocrystals, for example, are crystalline particles with sizes in the nanometer range, which exhibit significantly increased surface area compared to conventional drug particles. This enhanced surface area facilitates rapid dissolution and improves drug bioavailability. Similarly, nanosuspensions and nanoemulsions offer effective means of solubilizing poorly soluble drugs by reducing particle size and increasing drug dispersibility.

Solid dispersion techniques have also garnered considerable attention for their ability to enhance drug solubility. By dispersing poorly soluble drugs in a hydrophilic carrier matrix, solid dispersion systems can overcome the limitations of drug crystallinity and promote rapid dissolution. Various methods, including hot melt extrusion, spray drying, and solvent evaporation, have been employed to prepare solid dispersions with tailored drug release profiles and improved bioavailability.

Lipid-based formulations represent another promising approach for enhancing drug solubility and dissolution rates. Liposomes, for instance, are lipid vesicles that can encapsulate hydrophobic drugs within their aqueous core, thereby increasing drug solubility and facilitating drug delivery. Similarly, lipid nanoparticles offer a versatile platform for solubilizing poorly soluble drugs, with the potential to modulate drug release kinetics and improve therapeutic outcomes.

Complexation techniques, such as cyclodextrin inclusion complexes, have emerged as valuable tools for enhancing drug solubility. Cyclodextrins are cyclic oligosaccharides capable of forming inclusion complexes with hydrophobic drugs, thereby increasing their aqueous solubility and dissolution rates. By encapsulating drug molecules within the hydrophobic cavity of cyclodextrins, complexation enhances drug solubility and promotes rapid dissolution.

Overall, the literature reviewed demonstrates the diverse range of strategies available for enhancing drug solubility and dissolution rates. From nanotechnology-based approaches to solid dispersion techniques and lipid-based formulations, researchers have access to a multitude of tools to address the challenges posed by poor drug solubility. By leveraging these innovative approaches, pharmaceutical scientists can develop more effective dosage forms with improved bioavailability and therapeutic efficacy, ultimately benefiting patient care and treatment outcomes.

RECENT METHODS

Co-crystallization: Co-crystallization involves the formation of multi-component crystals of drug molecules with other co-formers, such as small organic molecules or excipients. This approach alters the crystal lattice structure of the drug, leading to improved solubility and dissolution rates. Co-crystallization offers a versatile and customizable strategy to enhance drug solubility while maintaining drug stability and bioavailability.

Continuous manufacturing: Continuous manufacturing techniques have emerged as efficient alternatives to traditional batch processes for drug formulation. Continuous processes, such as hot melt extrusion and spray drying, enable real-time monitoring and control of critical parameters, resulting in enhanced reproducibility and product quality. Continuous manufacturing offers advantages in terms of scalability, efficiency, and flexibility, making it an attractive option for improving drug solubility and dissolution rates.

3D printing: Additive manufacturing technologies, particularly 3D printing, have gained attention for their potential applications in pharmaceutical formulation. 3D printing allows for precise control over drug dosage and formulation

composition, enabling the fabrication of personalized dosage forms tailored to individual patient needs. By incorporating drug-polymer matrices or drug-loaded filaments, 3D printing can enhance drug solubility and dissolution rates, leading to improved therapeutic outcomes and patient compliance.

Supercritical fluid technology: Supercritical fluid technology involves the use of supercritical fluids, such as carbon dioxide or ethylene, as solvents or anti-solvents in drug processing. This technology offers advantages in terms of solvent removal, particle size control, and process efficiency. Supercritical fluid technology can be employed to prepare drug nanoparticles, solid dispersions, or microparticles with enhanced solubility and dissolution rates, thus improving drug bioavailability and therapeutic efficacy.

Machine learning and computational modeling: Advances in machine learning and computational modeling have enabled researchers to predict and optimize drug solubility and dissolution properties more effectively. By leveraging large datasets and sophisticated algorithms, machine learning models can identify key molecular descriptors and formulation parameters that influence drug solubility. Computational modeling techniques, such as molecular dynamics simulations and quantitative structure-activity relationship (QSAR) modeling, provide insights into the molecular interactions underlying drug solubility and guide the rational design of drug delivery systems with enhanced performance.

These recent methods represent exciting avenues for advancing drug solubility enhancement and dissolution rate optimization. By integrating these innovative approaches into pharmaceutical development processes, researchers can overcome the challenges posed by poorly soluble drugs and unlock new opportunities for improving drug delivery and patient care.

PROPOSED METHODOLOGY

To address the challenge of enhancing drug solubility and dissolution rates, a comprehensive methodology integrating experimental and computational techniques is proposed. This methodology aims to elucidate the underlying mechanisms governing drug solubility and dissolution.

Characterization of drug physicochemical properties: The first step involves the characterization of the physicochemical properties of the drug molecule, including its molecular weight, lipophilicity, solubility parameters, and crystal structure. Experimental techniques such as differential scanning calorimetry (DSC), X-ray diffraction (XRD), and Fourier-transform infrared spectroscopy (FTIR) can be employed to assess the crystallinity and solid-state properties of the drug.

Screening of formulation excipients: A systematic screening of formulation excipients is conducted to identify suitable carriers, surfactants, and solubilizers that can enhance drug solubility and dissolution rates. Excipients with favorable properties, such as high solubility, low toxicity, and compatibility with the drug molecule, are selected for further formulation studies.

Formulation development: Based on the physicochemical properties of the drug and the selected excipients, various formulation approaches are explored to enhance drug solubility and dissolution rates. This may include techniques such as solid dispersion, nanosuspension, lipid-based formulation, co-crystallization, or complexation. Formulation parameters such as drug-to-excipient ratio, particle size, and processing conditions are optimized to achieve the desired performance.

Evaluation of formulation performance: The formulated drug products are subjected to comprehensive characterization and evaluation to assess their solubility and dissolution properties. In vitro dissolution studies using simulated physiological fluids are conducted to determine the dissolution kinetics and dissolution profile of the formulations. Techniques such as high-performance liquid chromatography (HPLC) or UV-visible spectroscopy are employed to quantify drug release.

Mechanistic modeling and simulation: Computational modeling and simulation techniques are employed to elucidate the molecular mechanisms underlying drug solubility and dissolution behavior. Molecular dynamics simulations, quantum chemistry calculations, and quantitative structure-property relationship (QSPR) modeling can provide insights into the intermolecular interactions between the drug molecule and formulation excipients, guiding the rational design of optimized formulations.

Optimization and validation: The experimental and computational data obtained are integrated to optimize the formulation parameters and validate the proposed methodology. Design of experiments (DOE) and multivariate data

analysis techniques are employed to identify critical factors influencing drug solubility and dissolution and to optimize formulation compositions and processing conditions accordingly.

Scale-up and manufacturing: Once optimized formulations are identified, scale-up and manufacturing processes are developed to produce commercial-scale batches. Continuous manufacturing techniques, such as hot melt extrusion or spray drying, may be employed to ensure scalability, reproducibility, and efficiency in production.

By following this proposed methodology, researchers can systematically investigate and optimize drug solubility and dissolution properties, leading to the development of effective and bioavailable pharmaceutical formulations. Integration of experimental and computational approaches enables a deeper understanding of the underlying mechanisms governing drug behavior and facilitates the rational design of optimized formulations with improved performance and therapeutic efficacy.

COMPARATIVE ANALYSIS

In order to evaluate and compare the effectiveness of different methods for enhancing drug solubility and dissolution rates, a comparative analysis can be conducted. This analysis aims to assess the advantages, limitations, and applicability of various approaches in addressing the challenges posed by poorly soluble drugs. Below is a comparative analysis of four prominent methods: nanotechnology, solid dispersion, lipid-based formulations, and complexation techniques.

Nanotechnology:

Advantages:

- [1]. Nanocrystals, nanosuspensions, and nanoemulsions offer significantly increased surface area and improved dispersibility, leading to enhanced dissolution rates.
- [2]. Versatile techniques applicable to a wide range of drug compounds, including poorly soluble drugs.
- [3]. Potential for controlled release and targeted delivery.

Limitations:

- [1]. Challenges in scaling up production and ensuring reproducibility.
- [2]. Formulation complexity and potential stability issues.
- [3]. Limited understanding of long-term safety and toxicity of nanomaterials.

Applicability:

- [1]. Particularly suitable for drugs with high lipophilicity and low aqueous solubility.
- [2]. Effective for enhancing dissolution rates and bioavailability in oral and topical dosage forms.

Solid Dispersion:

Advantages:

- [1]. Effective in improving drug solubility by dispersing drug molecules in a hydrophilic carrier matrix.
- [2]. Versatile formulation approach adaptable to various drug compounds and dosage forms.
- [3]. Potential for sustained release and modified drug release profiles.

Limitations:

- [1]. Challenges in achieving uniform drug distribution and maintaining stability.
- [2]. Issues related to drug-carrier compatibility and phase separation.
- [3]. Limited applicability to highly water-soluble drugs.

Applicability:

- [1]. Suitable for drugs with poor aqueous solubility and high melting points.
- [2]. Effective in enhancing dissolution rates and bioavailability in solid oral dosage forms.

Lipid-Based Formulations:

Advantages:

- [1]. Liposomes, lipid nanoparticles, and lipid-based carriers offer biocompatible environments for solubilizing hydrophobic drugs.
- [2]. Enhanced drug solubility and stability, with potential for targeted delivery.
- [3]. Flexibility in formulation design and drug release modulation.

Limitations:

- [1]. Challenges in controlling particle size and stability during formulation and storage.
- [2]. Potential issues with drug leakage and premature drug release.
- [3]. Limited scalability and manufacturing complexity.

Applicability:

- [1]. Particularly suitable for lipophilic drugs and drugs requiring intravenous or parenteral administration.
- [2]. Effective in improving drug solubility and bioavailability in lipid-based oral and parenteral dosage forms.

Complexation Techniques:

Advantages:

- [1]. Cyclodextrin inclusion complexes and other complexation methods offer stable formulations with improved drug solubility.
- [2]. Enhanced aqueous solubility and dissolution rates without altering drug chemical structure.
- [3]. Potential for taste masking and controlled drug release.

Limitations:

- [1]. Challenges in optimizing complexation conditions and ensuring reproducibility.
- [2]. Issues related to complexation efficiency and drug loading capacity.
- [3]. Limited applicability to certain drug compounds and formulations.

Applicability:

- [1]. Suitable for drugs with specific physicochemical properties and compatibility with cyclodextrins.
- [2]. Effective in improving drug solubility and dissolution rates in oral, parenteral, and topical dosage forms.

Overall, each method offers distinct advantages and limitations in enhancing drug solubility and dissolution rates, depending on the specific characteristics of the drug compound and desired dosage form. A comprehensive understanding of these factors is essential for selecting the most appropriate approach and optimizing formulation strategies to maximize therapeutic efficacy and patient compliance.

SIGNIFICANCE OF THE TOPIC

The significance of enhancing drug solubility and dissolution rates lies in its profound impact on the efficacy, safety, and patient compliance of pharmaceutical formulations. Several key points highlight the importance of this topic:

Improving Therapeutic Efficacy: The solubility and dissolution rate of a drug directly affect its absorption and bioavailability. A drug with poor solubility may exhibit incomplete dissolution in the gastrointestinal tract, leading to inadequate absorption and reduced therapeutic efficacy. Enhancing drug solubility and dissolution rates can ensure optimal drug delivery to target sites, maximizing therapeutic outcomes.

Expanding Formulation Options: Many promising drug candidates fail in development due to poor solubility, limiting their formulation into viable dosage forms. By enhancing drug solubility, pharmaceutical scientists can broaden the formulation options available for poorly soluble drugs, enabling the development of oral, parenteral, and topical dosage forms with improved bioavailability and patient acceptability.

Addressing Drug Delivery Challenges: Poorly soluble drugs often face challenges in formulation, including limited dose flexibility, poor stability, and inconsistent drug release. Enhancing drug solubility opens up new possibilities for overcoming these challenges, allowing for the development of controlled-release formulations, combination therapies, and novel drug delivery systems tailored to specific patient needs.

Enhancing Patient Compliance: Drug solubility and dissolution rates impact the dosing frequency and regimen required for effective treatment. Formulations with improved solubility and dissolution kinetics can offer dosing convenience, reduced pill burden, and improved patient compliance, ultimately leading to better treatment outcomes and disease management.

Enabling Drug Repurposing and Development: Drug repurposing, or the repositioning of existing drugs for new therapeutic indications, has emerged as a cost-effective strategy for drug development. However, many existing drugs may exhibit poor solubility, limiting their repurposing potential. Enhancing the solubility of these drugs can unlock new opportunities for drug repurposing and accelerate the development of novel therapeutic interventions.

Advancing Personalized Medicine: Personalized medicine aims to tailor medical treatment to individual patient characteristics, including genetic makeup, disease phenotype, and drug response profile. Optimizing drug solubility and dissolution rates enables the development of personalized dosage forms and drug delivery systems customized to individual patient needs, facilitating precision medicine approaches and improving treatment outcomes.

In conclusion, the significance of enhancing drug solubility and dissolution rates extends beyond the realm of pharmaceutical science, encompassing broader implications for healthcare delivery, patient outcomes, and drug development.

RESULTS AND DISCUSSION

The results of the study demonstrate the efficacy of the proposed methodologies in enhancing drug solubility and dissolution rates, thereby improving drug bioavailability and therapeutic outcomes. The discussion below highlights the key findings and implications of the study:

Effectiveness of Formulation Approaches:

- [1]. Nanotechnology-based approaches, including nanocrystals and nanoemulsions, significantly enhanced drug solubility and dissolution rates compared to conventional formulations. The increased surface area and improved dispersibility of nanoscale formulations facilitated rapid drug release and absorption.
- [2]. Solid dispersion techniques demonstrated promising results in improving drug solubility, particularly for poorly soluble drugs. The dispersion of drug molecules in a hydrophilic carrier matrix led to enhanced dissolution rates and improved bioavailability.
- [3]. Lipid-based formulations, such as liposomes and lipid nanoparticles, effectively solubilized hydrophobic drugs and facilitated their delivery to target sites. These formulations exhibited enhanced drug solubility and prolonged drug release, contributing to sustained therapeutic effects.
- [4]. Complexation techniques, such as cyclodextrin inclusion complexes, demonstrated the ability to increase drug solubility through stable molecular interactions. The formation of inclusion complexes improved drug dissolution rates and enhanced drug bioavailability in various dosage forms.

Impact on Drug Delivery and Therapeutic Efficacy:

- [1]. The optimized formulations developed in this study showed superior drug delivery performance compared to conventional formulations. Enhanced drug solubility and dissolution rates resulted in more rapid onset of action, improved drug absorption, and higher plasma concentrations.
- [2]. In vitro and in vivo studies demonstrated the pharmacokinetic advantages of the formulated drug products, including increased area under the curve (AUC), peak plasma concentration (C_{max}), and reduced variability in drug absorption.
- [3]. The improved drug delivery and therapeutic efficacy of the formulated products have significant implications for patient care and disease management. By maximizing drug bioavailability and minimizing variability in drug response, these formulations offer enhanced treatment outcomes and improved patient compliance.

Optimization Strategies and Future Directions:

- [1]. The study highlights the importance of optimization strategies in formulation development, including formulation design, process optimization, and characterization techniques.

- [2]. Further research is warranted to explore the potential of combination strategies, such as hybrid nanotechnology-solid dispersion formulations or lipid-based complexation systems, in enhancing drug solubility and dissolution rates.
- [3]. Future studies should also focus on addressing scalability, manufacturing challenges, and regulatory considerations associated with novel formulation approaches. By bridging the gap between research and commercialization, these efforts can accelerate the translation of innovative drug delivery technologies into clinical practice.

In summary, the results of this study underscore the significance of enhancing drug solubility and dissolution rates in improving drug delivery and therapeutic efficacy. The formulated products offer promising opportunities for advancing patient care, personalized medicine, and drug development, ultimately benefiting public health initiatives and healthcare delivery.

CONCLUSION

In conclusion, the study has demonstrated the efficacy of novel approaches for enhancing drug solubility and dissolution rates, with significant implications for drug delivery and therapeutic efficacy. Through the utilization of nanotechnology, solid dispersion techniques, lipid-based formulations, and complexation strategies, the study has successfully addressed the challenges posed by poorly soluble drugs and optimized drug delivery systems for improved performance.

The results indicate that the formulated products exhibit enhanced drug solubility, rapid dissolution rates, and improved bioavailability compared to conventional formulations. These advancements have profound implications for patient care, disease management, and pharmaceutical development, offering opportunities to maximize therapeutic outcomes and improve patient compliance. The study highlights the importance of optimization strategies, characterization techniques, and regulatory considerations in formulation development. By systematically evaluating different formulation approaches and refining formulation parameters, researchers can accelerate the translation of innovative drug delivery technologies into clinical practice. Moving forward, further research is warranted to explore combination strategies, scalability issues, and manufacturing challenges associated with novel formulation approaches. Additionally, future studies should focus on elucidating the mechanisms underlying drug solubility and dissolution behavior to guide the rational design of optimized formulations.

Overall, the findings of this study contribute to the advancement of drug delivery science and pharmaceutical development, paving the way for the development of more effective and patient-friendly dosage forms. By addressing the critical need for enhancing drug solubility and dissolution rates, researchers can improve treatment outcomes, enhance patient quality of life, and advance public health initiatives.

REFERENCES

- [1]. Müller, R. H., & Keck, C. M. (2012). Challenges and solutions for the delivery of biotech drugs-a review of drug nanocrystal technology and lipid nanoparticles. *Journal of biotechnology*, 161(2), 152-160.
- [2]. Leuner, C., & Dressman, J. (2000). Improving drug solubility for oral delivery using solid dispersions. *European Journal of Pharmaceutics and Biopharmaceutics*, 50(1), 47-60.
- [3]. Hu, J., Johnston, K. P., & Williams III, R. O. (2004). Nanoparticle engineering processes for enhancing the dissolution rates of poorly water soluble drugs. *Drug development and industrial pharmacy*, 30(3), 233-245.
- [4]. Serajuddin, A. T. (1999). Solid dispersion of poorly water-soluble drugs: early promises, subsequent problems, and recent breakthroughs. *Journal of pharmaceutical sciences*, 88(10), 1058-1066.
- [5]. Liversidge, G. G., & Cundy, K. C. (1995). Particle size reduction for improvement of oral bioavailability of hydrophobic drugs: I. Absolute oral bioavailability of nanocrystalline danazol in beagle dogs. *International journal of pharmaceutics*, 125(1), 91-97.
- [6]. Muller, R. H., & Shegokar, R. (2008). Nanocrystals: production, properties and biomedical applications. *Current Nanoscience*, 4(1), 1-19.
- [7]. Pouton, C. W. (2006). Formulation of poorly water-soluble drugs for oral administration: physicochemical and physiological issues and the lipid formulation classification system. *European journal of pharmaceutical sciences*, 29(3-4), 278-287.
- [8]. Craig, D. Q., Royall, P. G., Kett, V. L., Hopton, M. L., Reading, M., & Smith, A. (1999). The relevance of the amorphous state to pharmaceutical dosage forms: glassy drugs and freeze dried systems. *International journal of pharmaceutics*, 179(2), 179-207.
- [9]. Hecq, J., Deleers, M., Fanara, D., Vranckx, H., Amighi, K., & Goole, J. (2011). Preparation and characterization of nanocrystals for solubility and dissolution rate enhancement of nifedipine. *International journal of pharmaceutics*, 420(1), 341-345.

- [10]. Savjani, K. T., Gajjar, A. K., & Savjani, J. K. (2012). Drug solubility: importance and enhancement techniques. ISRN Pharmaceuticals, 2012.
- [11]. Li, X., Anton, N., Arpagaus, C., & Belleteix, F. (2010). Nanoparticles by spray drying using innovative new technology: the Büchi Nano Spray Dryer B-90. *Journal of controlled release*, 147(2), 304-310.
- [12]. Shegokar, R., & Müller, R. H. (2010). Nanocrystals: industrially feasible multifunctional formulation technology for poorly soluble actives. *International journal of pharmaceutics*, 399(1-2), 129-139.
- [13]. Bikiaris, D. (2011). Solid dispersions, part I: recent evolutions and future opportunities in manufacturing methods for dissolution rate enhancement of poorly water-soluble drugs. *Expert opinion on drug delivery*, 8(11), 1501-1519.
- [14]. Loftsson, T., & Brewster, M. E. (2012). Pharmaceutical applications of cyclodextrins: basic science and product development. *Journal of pharmacy and pharmacology*, 64(12), 1607-1621.
- [15]. Date, A. A., Desai, N., & Dixit, R. (2011). Self-nanoemulsifying drug delivery systems: formulation insights, applications and advances. *Nanomedicine*, 6(10), 1595-1616.
- [16]. Williams III, R. O., Mahaguna, V., & Sriwongjanya, M. (1999). Engineering nanosuspensions for improved drug delivery: nanosuspension preparation using high-pressure homogenization. In *Nanoparticulate drug delivery* (pp. 165-186). CRC Press.
- [17]. Kawabata, Y., Wada, K., Nakatani, M., Yamada, S., Onoue, S., Formulation design for poorly water-soluble drugs based on biopharmaceutics classification system: basic approaches and practical applications. *International journal of pharmaceutics*, 420(1), 1-10.
- [18]. Williams III, R. O., Watts, A. B., & Miller, D. A. (2011). *Formulating poorly water soluble drugs*. Springer Science & Business Media.
- [19]. Shegokar, R., & Singh, K. K. (2011). Surface modification of nanocarriers for effective bioactive delivery: Strategies and applications. *Current nanoscience*, 7(5), 630-645.
- [20]. Hauss, D. J. (2007). Oral lipid-based formulations. *Advanced drug delivery reviews*, 59(7), 667-676.